

# Computer Graphics (CS 543)

## Lecture 4(Part 2): Linear Algebra for Graphics (Points, Scalars, Vectors)

Prof Emmanuel Agu


*Computer Science Dept.  
Worcester Polytechnic Institute (WPI)*






# Points, Scalars and Vectors


- Points, vectors defined relative to a coordinate system
- Example: Point  $(5,4)$



# Vectors




- Magnitude
- Direction
- **NO** position
- Can be added, scaled, rotated
- CG vectors: 2, 3 or 4 dimensions




# Points

- Location in coordinate system
- Cannot add or scale
- Subtract 2 points = vector



# Vector-Point Relationship




- Diff. b/w 2 points = vector

$$\mathbf{v} = Q - P$$

- point + vector = point

$$\mathbf{v} + P = Q$$




# Vector Operations



- Define vectors  
 $\mathbf{a} = (a_1, a_2, a_3)$   
 $\mathbf{b} = (b_1, b_2, b_3)$

Then vector addition:

$$\mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2, a_3 + b_3)$$




# Vector Operations



- Define scalar,  $s$
- Scaling vector by a scalar


$$\mathbf{as} = (a_1s, a_2s, a_3s)$$



**Note** vector subtraction:

$$\mathbf{a} - \mathbf{b}$$

$$= (a_1 + (-b_1), a_2 + (-b_2), a_3 + (-b_3))$$





# Vector Operations: Examples

- Scaling vector by a scalar

$$\mathbf{a}s = (a_1s, a_2s, a_3s)$$

- Vector addition:

$$\mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2, a_3 + b_3)$$

- For example, if  $\mathbf{a}=(2,5,6)$  and  $\mathbf{b}=(-2,7,1)$  and  $s=6$ , then

$$\mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2, a_3 + b_3) = (0, 12, 7)$$

$$\mathbf{a}s = (a_1s, a_2s, a_3s) = (12, 30, 36)$$



# Affine Combination



- Given a vector

$$\mathbf{a} = (a_1, a_2, a_3, \dots, a_n)$$

$$a_1 + a_2 + \dots + a_n = 1$$

- Affine combination: Sum of all components = 1

- Convex affine = affine + no negative component

i.e

$$a_1, a_2, \dots, a_n = \textit{non-negative}$$



# Magnitude of a Vector

- Magnitude of  $\mathbf{a}$

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$$

- Normalizing a vector (unit vector)

$$\hat{\mathbf{a}} = \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{\text{vector}}{\text{magnitude}}$$

- Note magnitude of normalized vector = 1. i.e

$$\sqrt{a_1^2 + a_2^2 + \dots + a_n^2} = 1$$



# Magnitude of a Vector

- Example: if  $\mathbf{a} = (2, 5, 6)$

- Magnitude of  $\mathbf{a}$        $|\mathbf{a}| = \sqrt{2^2 + 5^2 + 6^2} = \sqrt{65}$


- Normalizing  $\mathbf{a}$

$$\hat{\mathbf{a}} = \left( \frac{2}{\sqrt{65}}, \frac{5}{\sqrt{65}}, \frac{6}{\sqrt{65}} \right)$$



# Convex Hull

- Smallest convex object containing  $P_1, P_2, \dots, P_n$
- Formed by “shrink wrapping” points





# Dot Product (Scalar product)

- Dot product,

$$d = \mathbf{a} \cdot \mathbf{b} = a_1 \cdot b_1 + a_2 \cdot b_2 + \dots + a_n \cdot b_n$$

- For example, if  $a=(2,3,1)$  and  $b=(0,4,-1)$

then

$$\begin{aligned} a \cdot b &= (2 \times 0) + (3 \times 4) + (1 \times -1) \\ &= 0 + 12 - 1 = 11 \end{aligned}$$



# Properties of Dot Products

- Symmetry (or commutative):

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$$

- Linearity:

$$(\mathbf{a} + \mathbf{c}) \cdot \mathbf{b} = \mathbf{a} \cdot \mathbf{b} + \mathbf{c} \cdot \mathbf{b}$$

- Homogeneity:


$$(s\mathbf{a}) \cdot \mathbf{b} = s(\mathbf{a} \cdot \mathbf{b})$$

- And

$$|\mathbf{b}^2| = \mathbf{b} \cdot \mathbf{b}$$



# Angle Between Two Vectors




$$\mathbf{b} = (|\mathbf{b}| \cos \phi_b, |\mathbf{b}| \sin \phi_b)$$


$$\mathbf{c} = (|\mathbf{c}| \cos \phi_c, |\mathbf{c}| \sin \phi_c)$$

$$\mathbf{b} \cdot \mathbf{c} = |\mathbf{b}| |\mathbf{c}| \cos \theta$$


Sign of  $\mathbf{b} \cdot \mathbf{c}$ :



$$\mathbf{b} \cdot \mathbf{c} > 0$$




$$\mathbf{b} \cdot \mathbf{c} = 0$$



$$\mathbf{b} \cdot \mathbf{c} < 0$$

# Angle Between Two Vectors



- Find the angle b/w the vectors  $\mathbf{b} = (3,4)$  and  $\mathbf{c} = (5,2)$





# Angle Between Two Vectors

- Find the angle b/w the vectors  $\mathbf{b} = (3,4)$  and  $\mathbf{c} = (5,2)$ 
  - $|\mathbf{b}| = 5$ ,  $|\mathbf{c}| = 5.385$

$$\hat{\mathbf{b}} = \left( \frac{3}{5}, \frac{4}{5} \right) \quad \hat{\mathbf{b}} \cdot \hat{\mathbf{c}} = 0.85422 = \cos \theta$$

$$\theta = 31.326^\circ$$

# Standard Unit Vectors




Define

$$\mathbf{i} = (1, 0, 0)$$

$$\mathbf{j} = (0, 1, 0)$$

$$\mathbf{k} = (0, 0, 1)$$



So that any vector,

$$\mathbf{v} = (a, b, c) = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$$



# Cross Product (Vector product)

If

$$\mathbf{a} = (a_x, a_y, a_z) \quad \mathbf{b} = (b_x, b_y, b_z)$$

Then

$$\mathbf{a} \times \mathbf{b} = (a_y b_z - a_z b_y) \mathbf{i} - (a_x b_z - a_z b_x) \mathbf{j} + (a_x b_y - a_y b_x) \mathbf{k}$$

Remember using determinant


$$\begin{vmatrix} i & j & k \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix}$$

**Note:**  $\mathbf{a} \times \mathbf{b}$  is perpendicular to  $\mathbf{a}$  and  $\mathbf{b}$

# Cross Product



**Note:**  $\mathbf{a} \times \mathbf{b}$  is perpendicular to both  $\mathbf{a}$  and  $\mathbf{b}$



# Cross Product



Calculate  $\mathbf{a} \times \mathbf{b}$  if  $\mathbf{a} = (3, 0, 2)$  and  $\mathbf{b} = (4, 1, 8)$

# Cross Product



Calculate  $\mathbf{a} \times \mathbf{b}$  if  $\mathbf{a} = (3, 0, 2)$  and  $\mathbf{b} = (4, 1, 8)$

$$\mathbf{a} \times \mathbf{b} = -2\mathbf{i} - 16\mathbf{j} + 3\mathbf{k}$$


# Normal for Triangle using Cross Product Method



plane  $\mathbf{n} \cdot (\mathbf{p} - \mathbf{p}_0) = 0$

$$\mathbf{n} = (\mathbf{p}_2 - \mathbf{p}_0) \times (\mathbf{p}_1 - \mathbf{p}_0)$$

normalize  $\mathbf{n} \leftarrow \mathbf{n} / |\mathbf{n}|$




Note that right-hand rule determines outward face



# Newell Method for Normal Vectors

- Problems with cross product method:
  - calculation difficult by hand, tedious
  - If 2 vectors almost parallel, cross product is small
  - Numerical inaccuracy may result



- Proposed by Martin Newell at Utah (teapot guy)
  - Uses formulae, suitable for computer
  - Compute during mesh generation
  - Robust!





# Newell Method Example


- Example: Find normal of polygon with vertices  $P_0 = (6,1,4)$ ,  $P_1=(7,0,9)$  and  $P_2 = (1,1,2)$

- Using simple cross product:

$$((7,0,9)-(6,1,4)) \times ((1,1,2)-(6,1,4)) = (2,-23,-5)$$

$P_1 - P_0$

$P_2 - P_0$





# Newell Method for Normal Vectors

- Formulae: Normal  $N = (m_x, m_y, m_z)$

$$m_x = \sum_{i=0}^{N-1} (y_i - y_{next(i)}) (z_i + z_{next(i)})$$

$$m_y = \sum_{i=0}^{N-1} (z_i - z_{next(i)}) (x_i + x_{next(i)})$$

$$m_z = \sum_{i=0}^{N-1} (x_i - x_{next(i)}) (y_i + y_{next(i)})$$

Using Newell method, for previous example plug in values result is same:  
Normal is (2, -23, -5)



# Finding Vector Reflected From a Surface


- $\mathbf{a}$  = original vector
- $\mathbf{n}$  = normal vector
- $\mathbf{r}$  = reflected vector
- $\mathbf{m}$  = projection of  $\mathbf{a}$  along  $\mathbf{n}$
- $\mathbf{e}$  = projection of  $\mathbf{a}$  orthogonal to  $\mathbf{n}$

**Note:**  $\theta_1 = \theta_2$

$$\mathbf{e} = \mathbf{a} - \mathbf{m}$$

$$\mathbf{r} = \mathbf{e} - \mathbf{m}$$


$$\Rightarrow \mathbf{r} = \mathbf{a} - 2\mathbf{m}$$





# Lines

- Consider all points of the form
  - $P(\alpha) = P_0 + \alpha \mathbf{d}$
  - **Line:** Set of all points that pass through  $P_0$  in direction of vector  $\mathbf{d}$



# Parametric Form



- Two-dimensional forms of a line


- **Explicit:**  $y = mx + h$
- **Implicit:**  $ax + by + c = 0$
- **Parametric:**

$$x(\alpha) = \alpha x_0 + (1-\alpha)x_1$$

$$y(\alpha) = \alpha y_0 + (1-\alpha)y_1$$

- Parametric form of line


- More robust and general than other forms
- Extends to curves and surfaces






# Convexity

- An object is *convex* iff for any two points in the object all points on the line segment between these points are also in the object



convex



not convex




# Curves and Surfaces

- **Curves:** 1-parameter **non-linear** functions of the form  $P(\alpha)$
- **Surfaces:** two-parameter functions  $P(\alpha, \beta)$ 
  - Linear functions give planes and polygons



$P(\alpha)$



$P(\alpha, \beta)$



# References

- Angel and Shreiner, Interactive Computer Graphics, 6<sup>th</sup> edition, Chapter 3
- Hill and Kelley, Computer Graphics using OpenGL, 3<sup>rd</sup> edition, Sections 4.2 - 4.4