



Computer Graphics (CS 543)

Lecture 6 (Part 3): Derivation of Perspective Projection Transformation

Prof Emmanuel Agu



*Computer Science Dept.
Worcester Polytechnic Institute (WPI)*





Perspective Projection



- Projection – map the object from 3D space to 2D screen



Perspective()
Frustum()




Perspective Projection: Classical



Based on similar triangles:

$$\frac{y'}{y} = \frac{N}{-z}$$


$$\Rightarrow y' = y \times \frac{N}{-z}$$



Perspective Projection: Classical

- So (x^*, y^*) projection of point, (x, y, z) unto near plane N is given as:

$$(x^*, y^*) = \left(x \frac{N}{-z}, y \frac{N}{-z} \right)$$



- Numerical example:



Q. Where on the viewplane does $P = (1, 0.5, -1.5)$ lie for a near plane at $N = 1$?

$$(x^*, y^*) = \left(x \frac{N}{-z}, y \frac{N}{-z} \right) = \left(1 \times \frac{1}{1.5}, 0.5 \times \frac{1}{1.5} \right) = (0.666, 0.333)$$



Pseudodepth

- Classical perspective projection projects (x,y) coordinates to (x^*, y^*) , drops z coordinates




- But we need z to find closest object (depth testing)!!!



Perspective Transformation

- **Perspective transformation** maps actual z distance of perspective view volume to range $[-1 \text{ to } 1]$ (**Pseudodepth**) for canonical view volume





We want perspective Transformation and NOT classical projection!!


Set scaling z
Pseudodepth = $az + b$
Next solve for a and b



Perspective Transformation

- We want to transform viewing frustum volume into canonical view volume








Perspective Transformation using Pseudodepth

$$(x^*, y^*, z^*) = \left(x \frac{N}{-z}, y \frac{N}{-z}, \frac{az + b}{-z} \right)$$

- Choose a, b so as z varies from **Near** to **Far**, pseudodepth varies from **-1** to **1** (canonical cube)

- Boundary conditions
 - $z^* = -1$ when $z = -N$
 - $z^* = 1$ when $z = -F$





Transformation of z : Solve for a and b

- Solving:

$$z^* = \frac{az + b}{-z}$$

- Use boundary conditions


- $z^* = -1$ when $z = -N$(1)

- $z^* = 1$ when $z = -F$(2)

- Set up simultaneous equations

$$-1 = \frac{-aN + b}{N} \Rightarrow -N = -aN + b \dots \dots (1)$$

$$1 = \frac{-aF + b}{F} \Rightarrow F = -aF + b \dots \dots (2)$$



Transformation of z : Solve for a and b

$$-N = -aN + b \dots \dots (1)$$

$$F = -aF + b \dots \dots (2)$$

- Multiply both sides of (1) by -1


$$N = aN - b \dots \dots (3)$$

- Add eqns (2) and (3)

$$F + N = aN - aF$$

$$\Rightarrow a = \frac{F + N}{N - F} = \frac{-(F + N)}{F - N} \dots \dots (4)$$

- Now put (4) back into (3)



Transformation of z : Solve for a and b

- Put solution for a back into eqn (3)

$$N = aN - b \dots \dots (3)$$


$$\Rightarrow N = \frac{-N(F + N)}{F - N} - b$$

$$\Rightarrow b = -N - \frac{-N(F + N)}{F - N}$$

$$\Rightarrow b = \frac{-N(F - N) - N(F + N)}{F - N} = \frac{-NF - N^2 - NF + N^2}{F - N} = \frac{-2NF}{F - N}$$

- So

$$a = \frac{-(F + N)}{F - N} \qquad b = \frac{-2FN}{F - N}$$



What does this mean?


- Original point z in original view volume, transformed into z^* in canonical view volume

$$z^* = \frac{az + b}{-z}$$

- where

$$a = \frac{-(F + N)}{F - N}$$

$$b = \frac{-2FN}{F - N}$$





Homogenous Coordinates

- Want to express projection transform as 4x4 matrix
- Previously, homogeneous coordinates of
 $P = (P_x, P_y, P_z) \Rightarrow (P_x, P_y, P_z, 1)$
- Introduce arbitrary scaling factor, w , so that
 $P = (wP_x, wP_y, wP_z, w)$ (**Note:** w is non-zero)
- For example, the point $P = (2, 4, 6)$ can be expressed as
 - $(2, 4, 6, 1)$
 - or $(4, 8, 12, 2)$ where $w=2$
 - or $(6, 12, 18, 3)$ where $w = 3$, or....
- To convert from homogeneous back to ordinary coordinates, first divide all four terms by w and discard 4th term



Perspective Projection Matrix

- Recall Perspective Transform

$$(x^*, y^*, z^*) = \left(x \frac{N}{-z}, y \frac{N}{-z}, \frac{az + b}{-z} \right)$$

- We have: $x^* = x \frac{N}{-z}$ $y^* = y \frac{N}{-z}$ $z^* = \frac{az + b}{-z}$

- In matrix form:

$$\begin{pmatrix} N & 0 & 0 & 0 \\ 0 & N & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & -1 & 0 \end{pmatrix} \begin{pmatrix} wx \\ wy \\ wz \\ w \end{pmatrix} = \begin{pmatrix} wNx \\ wNy \\ w(az + b) \\ -wz \end{pmatrix} \Rightarrow \begin{pmatrix} x \frac{N}{-z} \\ y \frac{N}{-z} \\ \frac{az + b}{-z} \\ 1 \end{pmatrix}$$

Perspective Transform Matrix **Original vertex** **Transformed Vertex** **Transformed Vertex after dividing by 4th term**




Perspective Projection Matrix

$$\begin{pmatrix} N & 0 & 0 & 0 \\ 0 & N & 0 & 0 \\ 0 & 0 & a & b \\ 0 & 0 & -1 & 0 \end{pmatrix} \begin{pmatrix} wP_x \\ wP_y \\ wP_z \\ w \end{pmatrix} = \begin{pmatrix} wNP_x \\ wNP_y \\ w(aP_z + b) \\ -wP_z \end{pmatrix} \Rightarrow \begin{pmatrix} x \frac{N}{-z} \\ y \frac{N}{-z} \\ \frac{az + b}{-z} \\ 1 \end{pmatrix}$$


$$a = \frac{-(F + N)}{F - N} \quad b = \frac{-2FN}{F - N}$$


- In perspective transform matrix, already solved for a and b :
- So, we have transform matrix to transform \mathbf{z} values



Perspective Projection

- Not done yet!! Can now transform z!
- Also need to transform the $\mathbf{x} = (\text{left}, \text{right})$ and $\mathbf{y} = (\text{bottom}, \text{top})$ ranges of viewing frustum to $[-1, 1]$
- Similar to glOrtho, we need to translate and scale previous matrix along x and y to get final projection transform matrix
- we translate by
 - $-(\text{right} + \text{left})/2$ in x
 - $-(\text{top} + \text{bottom})/2$ in y
- Scale by:
 - $2/(\text{right} - \text{left})$ in x
 - $2/(\text{top} - \text{bottom})$ in y







Perspective Projection

- Translate along x and y to line up center with origin of CVV
 - $-(right + left)/2$ in x
 - $-(top + bottom)/2$ in y
- Multiply by translation matrix:

$$\begin{pmatrix} 1 & 0 & 0 & -(right + left) / 2 \\ 0 & 1 & 0 & -(top + bottom) / 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

**Line up centers
Along x and y**






Perspective Projection

- To bring view volume size down to size of of CVV, scale by
 - $2/(\text{right} - \text{left})$ in x
 - $2/(\text{top} - \text{bottom})$ in y
- Multiply by scale matrix:

$$\begin{pmatrix} \frac{2}{\text{right} - \text{left}} & 0 & 0 & 0 \\ 0 & \frac{2}{\text{top} - \text{bottom}} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

**Scale size down
along x and y**



Perspective Projection Matrix



Previous
Perspective
Transform
Matrix

$$\begin{matrix}
 \text{Scale} & & \text{Translate} & & \text{Previous} \\
 & & & & \text{Perspective} \\
 & & & & \text{Transform} \\
 & & & & \text{Matrix}
 \end{matrix}
 \left(\begin{array}{cccc}
 \frac{2}{right - left} & 0 & 0 & 0 \\
 0 & \frac{2}{top - bottom} & 0 & 0 \\
 0 & 0 & 1 & 0 \\
 0 & 0 & 0 & 1
 \end{array} \right) \times \left(\begin{array}{cccc}
 1 & 0 & 0 & -(right + left) / 2 \\
 0 & 1 & 0 & -(top + bottom) / 2 \\
 0 & 0 & 1 & 0 \\
 0 & 0 & 0 & 1
 \end{array} \right) \times \left(\begin{array}{cccc}
 N & 0 & 0 & 0 \\
 0 & N & 0 & 0 \\
 0 & 0 & a & b \\
 0 & 0 & -1 & 0
 \end{array} \right)$$

→

$$\left(\begin{array}{cccc}
 \frac{2N}{x \max - x \min} & 0 & \frac{right + left}{right - left} & 0 \\
 0 & \frac{2N}{top - bottom} & \frac{top + bottom}{top - bottom} & 0 \\
 0 & 0 & \frac{-(F + N)}{F - N} & \frac{-2FN}{F - N} \\
 0 & 0 & -1 & 0
 \end{array} \right)$$



**Final Perspective
Transform Matrix**

glFrustum(left, right, bottom, top, N, F) N = near plane, F = far plane

Perspective Transformation




- After perspective transformation, viewing frustum volume is transformed into canonical view volume




Geometric Nature of Perspective Transform



- a) Lines through eye map into lines parallel to z axis after transform
- b) Lines perpendicular to z axis map to lines perp to z axis after transform



Normalization Transformation





References

- Interactive Computer Graphics (6th edition), Angel and Shreiner
- Computer Graphics using OpenGL (3rd edition), Hill and Kelley