


IMGD 2905


Descriptive Statistics

Chapter 3



Summarizing Data

- With lots of playtesting, there is a lot of data
 - This is a good thing!
- But raw data is often just a pile of numbers
 - Rarely of interest
 - Or even sensible
- **Q:** How to summarize all this information?



Summarizing Data


- With lots of playtesting, there is a lot of data
 - This is a good thing!
- But raw data is often just a pile of numbers
 - Rarely of interest
 - Or even sensible
- **Q:** How to summarize all this information?



Measures of **central tendency**

Examples? **Pros** and **Cons**?

Breakout 2



4 3 7 8 3 4 22 3 5 3 2 3

- Different for *central tendency* with **one** number?
- What are *pros* and *cons* of each?
- Icebreaker, Groupwork, Questions

<https://web.cs.wpi.edu/~imgd2905/d20/breakout/breakout-2.html>

Measure of Central Tendency: Mean

The sum of the measurements

divided by the number of
measurements

$$(6 + 4 + 5 + 4 + 8 + 3) / 6 = 5.$$

gives you the
mean.

<http://www.cdn.sciencebuddies.org/Files/463/9/MeanEquation.jpg>


- Also called the “**arithmetic mean**” or “**average**”
- In Excel, =AVERAGE(range)
=AVERAGEIF() – averages if numbers meet certain condition



Measure of Central Tendency: Median

- Sort values low to high and take middle value

Median



<https://betterexplained.com/wp-content/uploads/average/median.png>

10 11 13 15 16 23 26

middle number

<https://www.mathsisfun.com/definitions/images/median.gif>

13 22 26 38 36 42 49 50 77 81 98 110

Median = 45.5

<http://www.nedarc.org/statisticalHelp/basicStatistics/measuresOfCenter/images/median.gif>


- In Excel, =MEDIAN(range)



Measure of Central Tendency: Mode






- Number which occurs most frequently
 - Not too useful in many cases
- Best use for **categorical data**
- e.g., most popular Hero group in Heroes of the Storm

- In Excel, =MODE ()








<http://pad3.whstatic.com/images/thumb/c/cd/Find-the-Mode-of-a-Set-of-Numbers-Step-7.jpg/aid130521-v4-728px-Find-the-Mode-of-a-Set-of-Numbers-Step-7.jpg>

Depiction: Mean, Median, Mode?



Depiction: Mean, Median, Mode?




Which to Use, Mean, Median, Mode?

Which to Use, Mean, Median, Mode?

- **Mean** many statistical tests with sample
 - Estimator of population mean
 - Uses all data
- **Median** is useful for skewed data
 - e.g., income data (US Census) or housing prices (Zillow)
 - e.g., *Overwatch* team (6 players): 5 people level 5, 1 person level 275
 - Mean is 50 - not so useful since no one at this level
 - Median is 5 - more representative
 - Does not use all data. “Resistant” to extremes (e.g., 275)
 - But what if were exam scores? Hard to “bring up” grade
- **Mode** is useful primarily for categorical data only
 - Most played League champion, most popular maze, ...


Other Measures of Position

- May not always want center
 - e.g., want to know best LoL Champions
- What other positions may be desired?





Other Measures of Position

- May not always want center
 - e.g., want to know best LoL Champions
- Maximum / Minimum
 - Not discussed more
- Trimmed Mean
- Quartiles
- Percentiles



Trimmed Mean


- Take “trimming” off top and bottom (typically 5% or 10%)
 - Reduces effects of extreme values, like median
- In Excel, =TRIMMEAN(array, percent) 




Blue – original mean
Red – trimmed mean

Quartiles

- Sort values
- First quartile (**Q1**) is 25% from bottom
- Third quartile (**Q3**) is 75% from bottom
- (What is second quartile?)
- In Excel, =QUARTILE(array,n)




<https://www.hackmath.net/images/quartiles.png>




<https://mathbitsnotebook.com/Algebra1/StatisticsData/quartileboxview2.png>

Percentiles


- Generalization of quartiles
- N^{th} percentile is data point $n\%$ from bottom of data
- Interpolate as for first quartile
- In Excel, =PERCENTILE(array, k) (k: 0 to 1)



<https://www.mathsisfun.com/data/images/percentile-80.svg>




<http://www.psychometric-success.com/images/AA1301.gif>



http://www.isical.ac.in/~jeexiiscore_normal/PercentilesAdvantages.htm

Summarizing Data, Part 2

- Ok, pile of numbers can now be summarized as *one* number
 - Mean, median, mode
- But is that enough?
- Q: What other major aspect of numbers haven't we summarized?



Summarizing Data, Part 2

- Ok, pile of numbers can now be summarized as *one* number
 - Mean, median, mode
- But is that enough?
- Q: What other major aspect of numbers haven't we summarized?





Measures of **variation**
(aka measures of *dispersion*, or
measures of *spread*)

Summarizing Data, Part 2

“Then there is the man who drowned crossing a stream with an average depth of six inches.” – W.I.E. Gates



- Summarizing by single number rarely enough → need statement about **dispersion** (aka variation)



Above: does single number (**mean**) tell you enough about data?


Dispersion Overview (1 of 3)

- Is data clumped or spread out?




Dispersion Overview (2 of 3)

- Is data clumped or spread out?






Dispersion Overview (3 of 3)

- Is data clumped or spread out?




[“Motion and Scene Complexity for Streaming Video Games”](#)

What are Some Measures of Dispersion?



Breakout 3



Set A: 2 4 6 8 10

Set B: 2 9 9 10 10

- Different ways to report *dispersion* with **one** number?
- What are **pros** and **cons** of each?
- Icebreaker, Groupwork, Questions

<https://web.cs.wpi.edu/~imgd2905/d20/breakout/breakout-3.html>

Range

- Difference between smallest and largest value
- Somewhat obvious, but doesn't tell you much about "clumping"
 - Minimum may be zero
 - Maximum can be from outlier
 - Event not related to phenomena studied (e.g., 0 on project)
 - Maximum gets larger with # samples, so no "stable" point




In Excel, =MAX(array) - MIN(array)

12, 25, 27, 29, 36, 38, 40, 43, 50, 54, 62

Range = 62 - 12 = 50

<http://idolosol.com/images/range-3.jpg>



Variance

- Compute **mean** of sample
- Compute how far each value in sample is from **mean**
 - Some can be less than **mean**, some greater
 - So square this difference (why square?)
- Divide by number of sample values – 1
 - The “-1” corrects “bias” when trying to estimate *population variance* using *sample variance*

$$\text{Sample Variance} = s^2 = \frac{\overset{\text{“sum up all”}}{\Sigma} (X - \overset{\text{“mean”}}{\bar{X}})^2}{n - 1}$$

Variance Example

- Sample kills in *League of Legends* match
 - 12, 20, 16, 18, 19
 - What is sample variance?

- First, **mean** = $85 / 5 = 17$

<u>Kills</u>	<u>$X - \text{mean}$</u>	<u>$(X - \text{mean})^2$</u>
12	-5	25
20	3	9
16	-1	1
18	1	1
19	2	4

$$s^2 = (25 + 9 + 1 + 1 + 4) / (5 - 1) = 40 / 4 = 10 \text{ kills squared}$$

- In Excel, =VAR(array)




“Larger” means
“more spread”
... but units odd

Standard Deviation

- Square-root of variance
- Usually, use standard deviation instead of variance
 - Why? → Same *units* as data (e.g., “kills” in previous example)
- Can compare standard deviation to mean (*coefficient of variation*, next)
- But first:
 - Mendenhall’s Empirical Rule
 - Z-score


$$s = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

Low Standard Deviation




A “thin” curve means that your winrates remain close to the mean average.

High Standard Deviation




A “fat” curve means that there is a wider spread of your winrates.



Mendenhall's Empirical Rule

1. About **68% data** within one standard deviation of **mean**
 - interval between **mean-s** and **mean+s** contains about 68% of data
2. About **95%** within 2 standard deviations of mean
3. **Almost all** data within 3 standard deviations of mean




<https://mathbitsnotebook.com/Algebra1/StatisticsData/normalgrapha.jpg>

Rule assumes normal ("Bell curve") distribution

Z-Score

- Measure of how “far” from center (**mean**) single data point is
 - *Not* measure of dispersion for whole data set

$$Z = \frac{X - \bar{X}}{S}$$



<https://www.animatedsoftware.com/pics/stats/sgzscor2.gif>

Example

Mean	469
Std dev	119
X	650


Z-score for X ?
(650 – 469)/119 **1.52**

Coefficient of Variation (CV)

- Size of **standard deviation** relative to **mean**
 - e.g., large sd & large mean, not so spread
 - but large sd & small mean, more spread
- **Standard deviation** divided by **mean**
 - Can do this since same units!
- CV is “unit-less”, so measure of spread independent of quantity
 - E.g. seconds, clicks, spaces

Shown as percent (multiply by 100)


$$CV = \frac{s}{\bar{x}} \times 100$$




http://images.slideplayer.com/35/10391754/slides/slide_59.jpg

What is the relative CV for each curve?


<http://goo.gl/wrfVtH>



Same Means
Different Standard Deviations




Different Means
Same Standard Deviations



Different Means
Different Standard Deviations

Semi-Interquartile Range

- $\frac{1}{2}$ distance between **Q3** (75th percentile) and **Q1** (25th percentile)



<http://www.bbc.co.uk/staticarchive/9629000486ef4b1a40efa565c162cb779e0bd82c.png>

$$\frac{\text{Q3} - \text{Q1}}{2}$$

- Guideline: use semi-interquartile (SIQR) for index of dispersion whenever using **median** as index of central tendency

Index of Dispersion Example

(sorted)
Lap Times

1.9

2.7

3.9

4.1

4.2

4.2

4.4

4.5

4.5

4.8

4.9

5.1

5.1

5.3



5.6

5.9

- First, sort. Then, compute:
 - Mean = 4.4
 - Min = 1.9, Max = 5.9
 - Median = $[16 / 2] = 8^{\text{th}} = 4.5$
 - Q1 = $16 / 4 = 8^{\text{th}} = 4.1$
 - Q3 = $3 * 16 / 4 = 12^{\text{th}} = 5.1$
- $SIQR = (Q3 - Q1) / 2 = 0.5$
- $Variance = 0.96$
- $Stddev = 0.98$
- $CV = \text{stddev} / \text{mean} = 0.22$
- $Range = \text{max} - \text{min} = 4$

Breakout 4

- Group of 3!
- Rank *measures of dispersion* by sensitivity) to outliers
 - Variance
 - Range
 - Standard Deviation
 - Coefficient of Variation
 - Semi-interquartile Range



<https://web.cs.wpi.edu/~imgd2905/d20/breakout/breakout-4.html>


Ranking of Affect by Outliers?

Measure of Dispersion

- Variance
- Range
- Standard Deviation
- Coefficient of Variation
- Semi-interquartile Range

Most to Least

?



<http://www.a-levelmathstutor.com/images/statistics/outliers-graph01.jpg>


Ranking of Affect by Outliers?

Measure of Dispersion

- Variance
- Range
- Standard Deviation
- Coefficient of Variation
- Semi-interquartile Range

Most to Least

- Range **susceptible**
- Variance
 - Standard Deviation
 - Coefficient of Variation
- SIQR **resistant**



<http://www.a-levelmathstutor.com/images/statistics/outliers-graph01.jpg>

Only for **quantitative** data!
categorical can't quantify spread
since no 'distance'
Instead, give categories for given
percentile of samples
e.g., "90% of samples are in
3 categories"

Depicting Dispersion in Charts


- Histogram (next unit)
- Cumulative distribution (next unit)
- Box-and-Whiskers
- Error Bars

Box-and-Whiskers Chart


- Way of showing variation
- Highlight middle 50% (interquartile range, IQR)
 - “Box”
- Lines go to smallest non-outlier
 - “Whiskers”
- Points indicate outliers
- Middle line shows median
- Sometimes with mean
- **Outlier?** → Data value “way out there”, “far” from the rest
 - Formally, 1.5+ IQRs away from quartile
- Available in Excel



Also called “boxplot”



<http://support.sas.com/documentation/cdl/en/vaugh/65747/HTML/default/images/boxplot.png>



<https://support.office.com/en-us/article/Create-a-box-and-whisker-chart-62f4219f-db4b-4754-aca8-4743f6190f0d>

Error Bars for Columns and Points


- Line through graph point parallel to axis with “caps”
- Denotes uncertainty (variation) in value



Excel: click “+” → “Error Bars” → “type”

- Often:
 - 1 standard deviation
- Can be (discuss later):
 - 1 standard error
 - 1 confidence interval

State clearly!



<https://s3.amazonaws.com/cdn.graphpad.com/faq/804/images/804b.jpg>



<http://www.excel-easy.com/examples/images/error-bars/error-bars.png>